



IceBridge BedMachine Greenland, Version 6

USER GUIDE

How to Cite These Data

As a condition of using these data, you must include a citation:

Morlighem, M., Williams, C., Rignot, E., An, L., Arndt, J. E., Bamber, J., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B., O'Cofaigh, C., Palmer, S. J., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., & Zinglensen, K. (2025). *IceBridge BedMachine Greenland (IDBMG4, Version 6)* [Data set]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center.
<https://doi.org/10.5067/6B6B225B8V2D> [Date Accessed].

Literature Citation

As a condition of using these data, we request that you acknowledge the authors of this data set by referencing the following peer-reviewed publication.

Morlighem, M., Williams, C., Rignot, E., An, L., Arndt, J. E., Bamber, J., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B., O'Cofaigh, C., Palmer, S. J., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., & Zinglensen, K. (2017). BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multi-beam echo sounding combined with mass conservation. *Geophysical Research Letters*, 44. <https://doi.org/10.1002/2017GL074954>

FOR QUESTIONS ABOUT THESE DATA, CONTACT NSIDC@NSIDC.ORG

FOR CURRENT INFORMATION, VISIT <https://nsidc.org/data/IDBMG4>



National Snow and Ice Data Center

TABLE OF CONTENTS

1	DATA DESCRIPTION	2
1.1	Summary	2
1.2	File Information.....	2
1.2.1	Format.....	2
1.2.2	File Contents.....	2
1.2.3	Naming Convention	3
1.2.4	Final Map Images	3
1.3	Spatial Information.....	3
1.3.1	Coverage	3
1.3.2	Resolution.....	4
1.3.3	Geolocation.....	4
1.4	Temporal Information	4
1.4.1	Coverage	4
1.4.2	Resolution.....	4
2	DATA ACQUISITION AND PROCESSING.....	4
2.1	Background	4
2.2	Acquisition	5
2.3	Processing.....	7
2.4	Quality, Errors, and Limitations	7
3	VERSION HISTORY	9
4	ACKNOWLEDGMENTS	10
5	REFERENCES	10
6	DOCUMENT INFORMATION.....	16
6.1	Publication Date	16
6.2	Date Last Updated.....	16

1 DATA DESCRIPTION

1.1 Summary

This data set contains a bed topography and bathymetry map of Greenland based on mass conservation, multi-beam data, and other techniques. It also includes surface elevation and ice thickness data, as well as an ice/ocean/land mask.

1.2 File Information

1.2.1 Format

The data are provided as a single netCDF (.nc) file. The bed elevation parameter is also provided as a geoTIFF file (.tif).

1.2.2 File Contents

All parameters of the data set are detailed in Table 1.

Table 1. File Parameter Descriptions

Parameter	Description
bed	Bed elevation (bed topography) relative to the geoid (m)
dataid	Input data source ID: 1 = gimpdem, 2 = radar, 7 = seismic bathymetry, 10 = multibeam bathymetry
errbed	Bed topography/ice thickness error (m)
geoid	Height difference between the EIGEN-6C4 Geoid and WGS84 Ellipsoid (m)
mapping	WGS 84 / NSIDC Sea Ice Polar Stereographic North
mask	Mask: 0 = ocean; 1 = ice-free land; 2 = grounded ice; 3 = floating ice
rgi	Randolph Glacier Inventory (RGI) mask v7.0; 1 = inside RGI and 0 = outside RGI
source	Output data source ID: 0 = none; 1 = gimpdem; 2 = mass conservation; 3 = synthetic; 4 = interpolation; 5 = hydrostatic equilibrium; 6 = kriging; 7 = RTOPO-2; 8 = gravity inversion; 9 = IceBoost; 10+ = bathymetry data
surface	Ice surface elevation relative to the geoid (m)
thickness	Ice thickness (m)
x	Projection x coordinate (m)
y	Projection y coordinate (m)

1.2.3 Naming Convention

The netCDF data file is named `BedMachineGreenland-v6.nc` where "v6" refers to the version of the data set. The geoTIFF file is named `BedMachineGreenland_bed-v6.tif`.

1.2.4 Final Map Images

Figure 1 illustrates Greenland bed topography (m) with respect to mean sea level (left) and ice thickness (m) (right).




Figure 1. Greenland bed topography and ice thickness

1.3 Spatial Information

1.3.1 Coverage

Spatial coverage for this data set includes Greenland and portions of the Arctic.

Southernmost Latitude: 59° N

Northernmost Latitude: 84° N

Westernmost Longitude: 90° W

Easternmost Longitude: 9° E

1.3.2 Resolution

The output product is generated at a 150 m resolution. The true resolution varies between 150 m and 5 km.

1.3.3 Geolocation

WGS84 / NSIDC Sea Ice Polar Stereographic North (EPSG 3413)

1.4 Temporal Information

1.4.1 Coverage

The data were collected between 1 January 1993 and 31 December 2021. The nominal year of this data set is 2007.

1.4.2 Resolution

Not specified

2 DATA ACQUISITION AND PROCESSING

2.1 Background

Bed topography and seafloor bathymetry regulate ice dynamics and water flow around Greenland, which impact glacier undercutting, calving, and movement. High-resolution mapping of subglacial features can improve understanding of ice-ocean interactions and ice sheet changes. Bed topography and ice thickness data are primarily collected by airborne radar sounders; however, the data quality is insufficient along coastal margins. Using multiple data sources and methods, along with the radar-derived ice thickness, optimizes mapping to provide a comprehensive bed topography and fjord bathymetry map of Greenland.

The Center for Remote Sensing of Ice Sheets (CReSIS) Multichannel Coherent Radar Depth Sounder (MCoRDS) operates over a 180 to 210 MHz frequency range with multiple receivers

developed for airborne sounding and imaging of ice sheets. See [IceBridge MCoRDS L2 Ice Thickness, Version 1](#) for further information on the MCoRDS radar and the Level-2 data.

2.2 Acquisition

Source data used in deriving this product are listed below.

Table 2. Source Data

Type	Source
Ice thickness data	Operation IceBridge radar-derived, posted at 30–60 m, with a vertical precision of 30 m, collected by the MCoRDS radar (IceBridge MCoRDS L2 Ice Thickness)
	Doppler focused radar of the Technical University of Denmark (DTU) for the region of 79 North (Thomsen et al., 1997; Christensen et al., 2000) and Russell (Lindbäck et al., 2014)
	High CApability Radar Sounder (HiCARS; Peters et al., 2005; Peters et al., 2007) operated by the University of Texas, Institute for Geophysics
	Pathfinder Advanced Radar Ice Sounder (PARIS; Raney, 2010)
	Alfred Wegener Institute (AWI; Nixdorf et al., 1999)
	Uppsala University (UU; Lindbäck et al., 2014), collected in the vicinity of Russell Gletscher
	Millan et al. (2022) for ice caps
	IceBoost for periphery glaciers (Maffezzoli et al., 2025)
Sounding data	Multibeam echo sounding data from Oceans Melting Greenland (OMG 2016, 2017, and 2020 Missions) along the coast of West and Southeast Greenland
Bathymetry data	Oceans Melting Greenland (OMG 2021 Mission)
	Slabon et al. (2016) along the Northwest coast
	Weinrebe et al. (2009) in Torssukataq and Uummannaq Fjords
	O’Cofaigh et al. (2013) in Uummannaq Fjord
	Dowdeswell et al. (2014), Rignot et al. (2015), Fried et al. (2015), and Rignot et al. (2016) in Uummannaq Fjord
	Schumann et al. (2012), Holland et al. (2008) and Straneo et al. (2012) in Illulisat Icefjord
	Mix et al. (2015) in front of Petermann Fjord and the adjacent Hall Basin
	Sutherland et al. (2014) near Kangerdlussuaq
	Dowdeswell et al. (2016) in Nordvestfjord
	Chauché et al. (2014) near Lille Gletscher
	Straneo et al. (2016) in Sermilik fjord
	Motyka et al. (2017) in Godthåbsfjord
	Stevens et al. (2016) in Sarqardleq fjord

	Kjeldsen et al. (2017) in Timmiarmiut Fjord, Heimdal Glacier and Skjoldungen Fjord
	Rysgaard et al. (2003) in Young Sound fjord
	Bendtsen et al. (2017) near Flade Isblink Ice Cap
	Sutherland and Pickart (2008) on the continental shelf along the Southeast coast (single beam)
	Single-beam Olex and crowd sourced data from fishing and recreational vessels (MaxSea)
	An et al. (2019) along the Northwest Greenland coast
	Millan et al. (2018) and An et al. (2019) in Southeast Greenland
	An et al. (2021) in Northeast Greenland
	Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, in front of the Northeast Ice Stream
	Danish Hydrographic Office
Radar data	Ice velocity measurements derived from satellite radar data collected during 2008–2009, posted at 150 m, with errors of 10 m yr ⁻¹ in speed and 1.5° in flow direction (Mouginot et al., 2017): • Japanese Advanced Land Observing System (ALOS) PALSAR • Canadian RADARSAT-1 SAR • German TerraSAR-X • European Envisat Advanced SAR (ASAR)
	Center for Remote Sensing of Ice Sheets (CReSIS) (2017)
	Hiawatha thickness (Kjaer et al., 2018)
	Programme for Monitoring of the Greenland Ice Sheet (PROMICE) (Søren et al., 2019)
	East Greenland Ice Core Project (EGRIP) thickness (Franke et al., 2020)
	Northeast Greenland Ice Stream (NEGIS) thickness (Jansen et al., 2020)
Ancillary data	Surface Mass Balance (SMB) averaged for the years 1961 to 1990 downscaled to 1 km with a precision between 7 percent and 20 percent in the ablation zone (Noël et al., 2016; data set available from the authors upon request)
	Ice thickening rates from altimetry data differencing between the years 2003 and 2006 (Khan et al., 2014)
	Surface elevation from the Greenland Mapping Project (GIMP) Digital Elevation Model (Howat et al., 2014)
	Ice and Ocean mask from the Greenland Mapping Project (GIMP) Digital Elevation Model (Howat et al., 2014)
	RTopo-2 (Schaffer et al., 2016)

2.3 Processing

Sparse, airborne radar-sounding-derived ice thickness data are combined with comprehensive, high-resolution ice motion derived from satellite interferometric synthetic aperture radar to calculate ice thickness based on mass conservation. The mass conservation method solves the mass conservation equation to derive ice thickness while minimizing the departure from the original radar-derived ice thickness data. Ice surface motion provides a physical basis for extrapolating sparse ice thickness data to larger areas with few or no data. The method works best in areas of fast flow, where errors in the flow direction are small and glaciers slide on the bed. In the interior regions, where errors in the flow direction are larger, kriging is used to interpolate ice thickness for the 1993–2016 data (Morlighem et al., 2014). Beginning with the 2017 data, streamline diffusion was used in the interior regions instead of kriging.

Ocean bathymetry is mapped by combining sparse bathymetry measurements from single and multibeam measurements and casts and [RTopo-2](#) (Schaffer et al., 2016).

The algorithm neglects ice motion by internal shear, which is an excellent approximation for fast-flowing glaciers ($>100 \text{ m yr}^{-1}$) (Morlighem et al., 2014).

The bed topography is derived by subtracting the ice thickness from the [Greenland Mapping Project \(GIMP\) Digital Elevation Model](#).

2.4 Quality, Errors, and Limitations

Sources of error include error in ice velocity direction and magnitude, error in surface mass balance, and ice thinning rates. In a trial setting with unusually dense radar sounding coverage, errors in the mass-conservation-inferred thickness were 36 m, only slightly higher than that of the original data. In areas less well constrained by radar-derived thickness data or constrained by only one track of data, for example, in south Greenland, errors may exceed 50 m (Morlighem et al., 2013).

No or very little data were available for some fjords, and uncertainty may be high ($>500 \text{ m}$).

An error estimate of the bed elevation and ice thickness is provided in the data set, illustrated in Figure 2.




Figure 2. Error Estimate of Greenland Bed Elevation and Ice Thickness

3 VERSION HISTORY

Table 3. Version History Summary

Version	Date	Description of Changes
5.0 (retire)	13 Jan 2026	Data access was removed for v5.0. Temporal coverage was 1 Jan 1993 to 31 Dec 2021.
6.0	11 Dec 2025	- Generated ensemble of bed topography maps using ICESat-2 time series and used the median depth instead of a single estimate for some glaciers - Used ice thickness estimates from IceBoost for periphery glaciers (Maffezzoli et al., 2025) - Added a new field 'RGI' that indicates which pixel belongs to a glacier in the Randolph Glacier Inventory - Added bathymetry data from the Danish Hydrographic Office - Updated bathymetry of Petermann based on Ciraci et al. (2023) gravity inversion
4.0 (retire)	19 Dec 2022	Data access was removed for v4.0. Temporal coverage was 1 Jan 1993 to 1 Oct 2020.
5.0	6 Sep 2022	Added bathymetry measurements from the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, in front of the Northeast Ice Stream, ice thickness for ice caps from Millan et al. (2022), and bathymetry data from the 2021 OMG campaign. A geoTIFF file of bed elevation was also added.
3.0 (retire)	27 May 2021	Data access was removed for v3.0. Temporal coverage was 1 Jan 1993 to 31 Dec 2016.
4.0	13 May 2021	Incorporated updated radar data to better constrain the ice thickness inland. Multibeam data for ocean bathymetry were updated to include data from 2017 and 2020 OMG campaigns. Bathymetry data were added from new gravity inversions. The format of surface, bed, and thickness was changed to "single". Streamline diffusion was implemented in the interior regions instead of kriging beginning January 2017.
2.0 (retire)	20 Apr 2020	Data access was removed for v2.0. Temporal coverage was 1 Jan 2007 to 31 Dec 2007.
1.0 (retire)	10 Sep 2019	Data access was removed for v1.0. Temporal coverage was 1 Jan 2007 to 31 Dec 2007.
3.0	25 Sep 2017	Added ocean bathymetry all around Greenland based on data from NASA's Ocean Melting Greenland (OMG) and other campaigns of bathymetry measurements. The subglacial bed topography was also updated by including more ice thickness data and constraining the ice thickness at the ice/ocean interface based on bathymetry data when available.

2.0	19 May 2015	Improved processing of some basins and added Operation IceBridge 2014 data. Heights were provided with respect to mean sea level, instead of the WGS84 ellipsoid. The geoid was included in an additional field in the data.
1.0	13 Jan 2015	Initial release

4 ACKNOWLEDGMENTS

We would like to thank Greenland Ice Sheet Ocean Research Coordination Network (GRISO RCN) for their help in finding available bathymetry data.

5 REFERENCES

An, L., Rignot, E., Millan, R., Tinto, K., & Willis, J. (2019). Bathymetry of northwest Greenland using “Ocean Melting Greenland” (OMG) high-resolution airborne gravity and other data. *Remote Sensing*, 11(131). <https://doi.org/10.3390/rs11020131>

An, L., Rignot, E., Chauche, N., Holland, D., Holland, D., Jakobsson, M., Kane, E., Wood, M., Klaucke, I., Morlighem, M., Velicogna, I., Weinrebe, W., & Willis, J. K. (2019). Bathymetry of southeast Greenland from oceans melting Greenland (OMG) data. *Geophysical Research Letters*, 46(20), 11197–11205. <https://doi.org/10.1029/2019GL083953>

An, L., Rignot, E., Wood, M., Willis, J. K., Mouginot, J., & Khan, S. A. (2021). Ocean melting of the Zachariae Isstrøm and Nioghalvfjærdsfjorden glaciers, northeast Greenland. *Proceedings of the National Academy of Sciences*, 118(2), e2015483118. <https://doi.org/10.1073/pnas.2015483118>

Bendtsen, J., Mortensen, J., Lennert, K., Ehn, J. K., Boone, W., Galindo, V., Hu, Y., Dmitrenko, I. A., Kirillov, S. A., Kjeldsen, K. K., Kristoffersen, Y., Barber, D. G., & Rysgaard, S. (2017). Sea ice breakup and marine melt of a retreating tidewater outlet glacier in northeast Greenland (81°N). *Scientific Reports*, 7(1), 4941. <https://doi.org/10.1038/s41598>

Chauché, N., Hubbard, A., Gascard, J.-C., Box, J. E., Bates, R., Koppes, M., Sole, A., Christoffersen, P., & Patton, H. (2014). Ice-ocean interaction and calving front morphology at two west Greenland tidewater outlet glaciers. *Cryosphere*, 8(4), 1457–1468. <https://doi.org/10.5194/tc-8-1457-2014>

Christensen, E. L., Reeh, N., Forsberg, R., Hjelm Jørgensen, J., Skou, N., & Woelders, K. (2000). A low-cost glacier-mapping system. *Journal of Glaciology*, 46(154), 531–537. <https://doi.org/10.3189/172756500781833142>

Ciraci, E., Rignot, E., Scheuchl, B., & Dini, L. (2023). Melt rates in the kilometer-size grounding zone of Petermann Glacier, Greenland, before and during a retreat. *Earth, Atmosphere, and Planetary Sciences*, 120(20). <https://doi.org/10.1073/pnas.2220924120>

CReSIS (2017). *Greenland Platform 3 Data*. Lawrence, Kansas, USA. Digital Media.
<http://data.cresis.ku.edu/>

Dowdeswell, J. A., Hogan, K. A., O'Cofaigh, C., Fugelli, E. M. G., Evans, J., & Noormets, R. (2014). Late Quaternary ice flow in a West Greenland fjord and cross-shelf trough system: submarine landforms from Rink Isbrae to Uummannaq shelf and slope. *Quaternary Science Review*, 92(SI), 292–309. <https://doi.org/10.1016/j.quascirev.2013.09.007>

Dowdeswell, J. A., Batchelor, C. L., Hogan, K. A., & Schenke, H. W. (2016). Nord-vestfjord: a major East Greenland fjord system. In *Atlas of Submarine Glacial Landforms: Modern, Quaternary and Ancient* (vol. 46, pp. 43–44). Geological Society. <https://doi.org/10.1144/M46.40>

Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Bamber, J. L., Box, J. E., & Bales, R. C. (2009). Higher surface mass balance of the Greenland Ice Sheet revealed by high-resolution climate modeling. *Geophysical Research Letters*, 36, 1–5.
<https://doi.org/10.1029/2009GL038110>

Franke, S., Jansen, D., Binder, T., Dörr, N., Helm, V., Paden, J., Steinhage, D., & Eisen, O. (2020). Bed topography and subglacial landforms in the onset region of the Northeast Greenland Ice Stream. *Annals of Glaciology*, 61(81), 143–153. <https://doi.org/10.1017/aog.2020.12>

Fried, M. J., Catania, G. A., Bartholomaeus, T. C., Duncan, D., Davis, M., Stearns, L. A., Nash, J., Shroyer, E., & Sutherland, D. (2015). Distributed subglacial discharge drives significant submarine melt at a Greenland tidewater glacier. *Geophysical Research Letters*, 42(21), 9328–9336.
<https://doi.org/10.1002/2015GL065806>

Holland, D., Thomas, R., De Young, B., Ribergaard, M., & Lyberth, B. (2008). Acceleration of Jakobshavn Isbrae triggered by warm subsurface ocean waters. *Nature Geoscience*, 1(10), 659–664. <https://doi.org/10.1038/ngeo316>

Howat, I. M., Negrete, A., & Smith, B. E. (2014). The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets. *The Cryosphere*, 8, 1509–1518.
<https://doi.org/10.5194/tc-8-1509-2014>

Jakobsson, M., et al. (2012). The International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0. *Geophysical Research Letters*, 39(12). <https://doi.org/10.1029/2012GL052219>

Jakobsson, M., Mayer, L. A., Bringenspar, C., Castro, C. F., Mohammad, R., Johnson, P., Ketter, T., Accettella, D., Amblas, D., An, L., Arndt, J. E., Canals, M., Casamor, J. L., Chauché, N., Coakley, B., Danielson, S., Demarte, M., Dickson, M., Dorschel, B., . . . Zinglarsen, K. B. (2020).

The international bathymetric chart of the arctic ocean version 4.0. *Scientific Data*, 7(1), 176.

<https://doi.org/10.1038/s41597-020-0520-9>

Jansen, D., Franke, S., Binder, T., Helm, V., Paden, J. D. (2020). Ice thickness from the northern catchment region of Greenland's 79 North Glacier, recorded with the airborne AWI UWB radar system. Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, PANGAEA. <https://doi.org/10.1594/PANGAEA.913193>

Khan, S. A., Kjaer, K. H., Bevis, M., Bamber, J. L., Wahr, J., Kjeldsen, K. K., Bjork, A. A., Korsgaard, N. J., Stearns, L. A., van den Broeke, M. R., Liu, L., Larsen, N. K., & Muresan, I. S. (2014). Sustained mass loss of the northeast Greenland ice sheet triggered by regional warming. *Nature Climate Change*, 4(4), 292–299. <https://doi.org/10.1038/nclimate2161>

Kjær, K. H., Larsen, N. K., Binder, T., Bjørk, A. A., Eisen, O., Fahnestock, M. A., Funder, S., Garde, A. A., Haack, H., Helm, V., Houmark-Nielsen, M., Kjeldsen, K. K., Khan, S. A., Machguth, H., McDonald, I., Morlighem, M., Mouginot, J., Paden, J. D., Waight, T. E., Weikusat, C., Willerslev, E., & MacGregor, J. A. (2018). A large impact crater beneath Hiawatha Glacier in northwest Greenland. *Science Advances*, 4(11). <https://doi.org/10.1126/sciadv.aar8173>

Kjeldsen, K. K., Weinrebe, W., Bendtsen, J., Bjørk, A. A., & Kjær, K. H. (2017). Multi-beam Bathymetry and CTD-Measurements in two fjord systems in Southeast Greenland. *Earth System Science Data Discussion*, 1–25. <https://doi.org/10.5194/essd-2017-29>

Lindbäck, K., Pettersson, R., Doyle, S. H., Helanow, C., Jansson, P., Kristensen, S. S., Stenseng, L., Forsberg, R., & Hubbard, A. L. (2014). High-resolution ice thickness and bed topography of a land-terminating section of the Greenland Ice Sheet. *Earth System Science Data*, 6, 331–338. <https://doi.org/10.5194/essd-6-331-2014>

Maffezzoli, N., Rignot, E., Barbante, C., Petersen, T., & Vascon, S. (2025). A gradient-boosted tree framework to model the ice thickness of the world's glaciers (IceBoost v1.1). *Geoscientific Model Development*, 18(9), 2545–2568. <https://doi.org/10.5194/gmd-18-2545-2025>

Millan, R., Rignot, E., Mouginot, J., Wood, M., Bjørk, A. A., & Morlighem, M. (2018). Vulnerability of southeast Greenland glaciers to warm Atlantic water from Operation IceBridge and Ocean Melting Greenland data. *Geophysical Research Letters*, 45(6), 2688–2696. <https://doi.org/10.1002/2017GL076561>

Millan, R., Mouginot, J., Rabatel, A., & Morlighem, M. (2022). Ice velocity and thickness of the world's glaciers. *Nature Geoscience*, 15, 124–129. <https://doi.org/10.1038/s41561-021-00885-z>

Mix, A. C., Jakobsson, M., & the Petermann-2015 Scientific Party (2015). Petermann-2015 Expedition Launches International Collaboration in Arctic Science. Tech. rep., Arctic Research Consortium of the United States.

Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., ... Zinglensen, K. B. (2017). BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multi-beam echo sounding combined with mass conservation. *Geophysical Research Letters*, *44*, <https://doi.org/10.1002/2017GL074954>

Morlighem, M., Rignot, E., & Willis, J. (2016b). Improving bed topography mapping of Greenland glaciers using NASA's Oceans Melting Greenland (OMG) data. *Oceanography*, *29*(4), 62–71. <https://doi.org/10.5670/oceanog.2016.99>

Morlighem, M., Rignot, E., Mouginot, J., Seroussi, H., & Larour, E. (2014). Deeply incised submarine glacial valleys beneath the Greenland Ice Sheet. *Nature Geoscience*, *7*, 418–422. <https://doi.org/10.1038/ngeo2167>

Morlighem, M., Rignot, E., Mouginot, J., Seroussi, H., & Larour, E. (2014). High-resolution ice-thickness mapping in South Greenland. *Annals of Glaciology*, *55*(67), 64–70. <https://doi.org/10.3189/2014AoG67A088>

Morlighem, M., Rignot, E., Mouginot, J., Wu, X., Seroussi, H., Larour, E., & Paden, J. (2013). High-resolution bed topography mapping of Russell Glacier, Greenland, inferred from Operation IceBridge data. *Journal of Glaciology*, *59*(218), 1015–1023. <https://doi.org/10.3189/2013JoG12J235>

Morlighem, M., Rignot, E., Seroussi, H., Larour, E., Ben Dhia, H., & Aubry, D. (2011). A mass conservation approach for mapping glacier ice thickness. *Geophysical Research Letters*, *38*(19), 1–6. <https://doi.org/10.1029/2011GL048659>

Motyka, R. J., Cassotto, R., Truffer, M., Kjeldsen, K. K., Van As, D., Korsgaard, N. J., Fahnestock, M., Howat, I., Langen, P. L., Mortensen, J., Lennert, K., & Rysgaard, S. (2017). Asynchronous behavior of outlet glaciers feeding Godthåbsfjord (Nuup Kangerlua) and the triggering of Narsap Sermia's retreat in SW Greenland. *Journal of Glaciology*, *63*(238), 288–308. <https://doi.org/10.1017/jog.2016.138>

Mouginot, J., Rignot, E., Scheuchl, B., & Millan, R. (2017). Comprehensive Annual Ice Sheet Velocity Mapping Using Landsat-8, Sentinel-1, and RADARSAT-2 Data. *Remote Sensing*, *9*(4), <https://doi.org/10.3390/rs9040364>

Nixdorf, U., Steinhage, D., Meyer, U., Hempel, L., Jenett, M., Wachs, P., & Miller, H. (1999). The newly developed airborne radio-echo sounding system of the AWI as a glaciological tool. *Annals of Glaciology*, *29*, 231–238. <https://doi.org/10.3189/172756499781821346>

Noël, B., van de Berg, W. J., Machguth, H., Lhermitte, S., Howat, I., Fettweis, X., & van den Broeke, M. R. (2016). A daily, 1 km resolution data set of downscaled Greenland ice sheet surface

mass balance (1958–2015). *The Cryosphere*, 10(5), 2361–2377. <https://doi.org/10.5194/tc-10-2361-2016>

O'Cofaigh, C., Dowdeswell, J. A., Jennings, A. E., Hogan, K. A., Kilfeather, A., Hiemstra, J. F., Noormets, R., Evans, J., McCarthy, D. J., Andrews, J. T., Lloyd, J. M., & Moros, M. (2013). An extensive and dynamic ice sheet on the West Greenland shelf during the last glacial cycle. *Geology*, 41(2), 219–222. <https://doi.org/10.1130/G33759.1>

OMG Mission (2020). *Bathymetry (sea floor depth) data from the ship-based bathymetry survey*. (OMG SDS; Version 1). <https://doi.org/10.5067/OMGEV-BTYSS>

Peters, M., Blankenship, D., & Morse, D. (2005). Analysis techniques for coherent air-borne radar sounding: Application to West Antarctic ice streams. *Journal of Geophysical Research Solid Earth*, 110(B6). <https://doi.org/10.1029/2004JB003222>

Peters, M. E., Blankenship, D. D., Carter, S. P., Kempf, S. D., Young, D. A., & Holt, J. W. (2007). Along-track focusing of airborne radar sounding data from West Antarctica for improving basal reflection analysis and layer detection. *IEEE Transactions on Geoscience and Remote Sensing*, 45(9), 2725–2736. <https://doi.org/10.1109/TGRS.2007.897416>

Raney, K. (2010). *IceBridge PARIS L2 ice thickness (IRPAR2; Version 1) [Data set]*. NSIDC. <https://doi.org/10.5067/OMEAKG6GIJNB>

Rignot, E., & Mouginot, J. (2012). Ice flow in Greenland for the International Polar Year 2008–2009. *Geophysical Research Letters*, 39(11). <https://doi.org/10.1029/2012GL051634>

Rignot, E., Fenty, I., Xu, Y., Cai, C., & Kemp, C. (2015). Undercutting of marine-terminating glaciers in West Greenland. *Geophysical Research Letters*, 42(14), 5909–5917. <https://doi.org/10.1002/2015GL064236>

Rignot, E., Fenty, I., Xu, Y., Cai, C., Velicogna, I., O'Cofaigh, C., Dowdeswell, J. A., Weinrebe, W., Catania, G., & Duncan, D. (2016). Bathymetry data reveal glaciers vulnerable to ice-ocean interaction in Uummannaq and Vaigat glacial fjords, west Greenland. *Geophysical Research Letters*, 43(6), 2667–2674. <https://doi.org/10.1002/2016GL067832>

Rysgaard, S., Vang, T., Stjernholm, M., Rasmussen, B., Windelin, A., & Kiilsholm, S. (2003). Physical conditions, carbon transport, and climate change impacts in a northeast Greenland fjord. *Arctic, Antarctic, and Alpine Research*, 35(3), 301–312. [https://doi.org/10.1657/1523-0430\(2003\)035\[0301:PCCTAC\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2003)035[0301:PCCTAC]2.0.CO;2)

Schaffer, J., Timmermann, R., Arndt, J. E., Kristensen, S. S., Mayer, C., Morlighem, M., & Steinhage, D. (2016). A global, high-resolution data set of ice sheet topography, cavity geometry, and ocean bathymetry. *Earth System Science Data*, 8(2), 543–557. <https://doi.org/10.5194/essd-8-543-2016>

- Schenk, T., & Csatho, B. (2012). A new methodology for detecting ice sheet surface elevation changes from laser altimetry data. *IEEE Transactions on Geoscience and Remote Sensing*, 50(9), 3302–3316. <https://doi.org/10.1109/TGRS.2011.2182357>
- Schumann, K., Voelker, D., & Weinrebe, W. R. (2012). Acoustic mapping of the Ilulissat Ice Fjord mouth, West Greenland. *Quaternary Science Reviews*, 40, 78–88. <https://doi.org/10.1016/j.quascirev.2012.02.016>
- Slabon, P., Dorschel, B., Jokat, W., Myklebust, R., Hebbeln, D., & Gebhardt, C. (2016). Greenland ice sheet retreat history in the northeast Baffin Bay based on high-resolution bathymetry. *Quaternary Science Reviews*, 154, 182–198. <http://doi.org/10.1016/j.quascirev.2016.10.022>
- Sørensen, L. S., Simonsen, S. B., Forsberg, R., Stenseng, L., Skourup, H., Kristensen, S. S., & Colgan, W. (2019). *Programme for monitoring of the Greenland ice sheet (PROMICE): Airborne survey*. Dataset published via Geological Survey of Denmark and Greenland and DTU-Space. <https://doi.org/10.22008/promice/data/airbornesurvey>
- Stevens, L. A., Straneo, F., Das, S. B., Plueddemann, A. J., Kukulya, A. L., & Morlighem, M. (2016). Linking glacially modified waters to catchment-scale subglacial discharge using autonomous underwater vehicle observations. *The Cryosphere*, 10(1), 417–432. <https://doi.org/10.5194/tc-10-417-2016>
- Straneo, F., Sutherland, D. A., Holland, D., Gladish, C., Hamilton, G. S., Johnson, H. L., Rignot, E., Xu, Y., & Koppes, M. (2012). Characteristics of ocean waters reaching Greenland's glaciers. *Annals of Glaciology*, 53(60), 202–210. <https://doi.org/10.3189/2012AoG60A059>
- Straneo, F., Hamilton, G. S., Stearns, L. A., & Sutherland, D. A. (2016). Connecting the Greenland Ice Sheet and the Ocean, a case study of Helheim Glacier and Sermilik Fjord. *Oceanography*, 29(4, SI), 34–45. <https://doi.org/10.5670/oceanog.2016.97>
- Sutherland, D. A., Straneo, F., & Pickart, R. S. (2014). Characteristics and dynamics of two major Greenland glacial fjords. *Journal of Geophysical Research Oceans*, 119(6), 3767–3791. <https://doi.org/10.1002/2013JC009786>
- Sutherland, D. A., & Pickart, R. S. (2008). The East Greenland Coastal Current: Structure, variability, and forcing. *Progress in Oceanography*, 78(1), 58–77. <https://doi.org/10.1016/j.pocean.2007.09.006>
- Thomsen, H. H., Reeh, N., Olesen, O. B., Bøggild, C. E., Starzer, W., Weidick, A., & Higgins, A. K. (1997). The Nioghalvfjærdsfjorden glacier project, North-East Greenland: a study of ice sheet response to climatic change. *Geology of Greenland Survey Bulletin*, 176, 95–103.
- Weinrebe, R. W., KuijpKui, A., Klauke, I., & Fink, M. (2009). Multibeam Bathymetry Surveys in Fjords and Coastal Areas of West-Greenland. *American Geophysical Union, Fall Meeting 2009*, 90(52). Abstract OS21A1152.

6 DOCUMENT INFORMATION

6.1 Publication Date

December 2025

6.2 Date Last Updated

January 2026